



# Use of nuclear receptor luciferase-based bioassays to detect endocrine active chemicals in a biosolids-biochar amended soil



Carolyn G. Anderson<sup>a</sup>, Geetika Joshi<sup>a</sup>, Daniel A. Bair<sup>a</sup>, Charlotte Oriol<sup>b</sup>, Guochun He<sup>c</sup>, Sanjai J. Parikh<sup>a,\*</sup>, Michael S. Denison<sup>c</sup>, Kate M. Scow<sup>a</sup>

<sup>a</sup> Department of Land, Air and Water Resources, University of California, Davis, One Shields Avenue, Davis, CA, 95616, USA


<sup>b</sup> Laboratoire HydroSciences Montpellier, Université Montpellier 2, Maison des Sciences de l'Eau, 300, Avenue du Professeur Emile Jeanbrau, Montpellier, 34095, France

<sup>c</sup> Department of Environmental Toxicology, University of California, Davis, One Shields Avenue, Davis, CA, 95616, USA

## HIGHLIGHTS

- Biosolids and biochar can have positive effects on plant growth.
- Some high temperature biochars may contain endocrine active chemicals.
- A soil-specific extraction and luciferase-based bioassay (CALUX) can be used to detect endocrine active chemicals in soil.

## GRAPHICAL ABSTRACT



## ARTICLE INFO

### Article history:

Received 22 December 2016

Received in revised form

14 March 2017

Accepted 8 April 2017

Available online 8 April 2017

Handling Editor: Frederic Leusch

### Keywords:

Biosolids

Biochar

Land application

Endocrine disruption

CALUX bioassay

## ABSTRACT

Biosolids are a potentially valuable source of carbon and nutrients for agricultural soils; however, potential unintended impacts on human health and the environment must be considered. Virtually all biosolids contain trace amounts endocrine-disrupting chemicals derived from human use of pharmaceuticals and personal care products (PPCPs). One potential way to reduce the bioavailability of PPCPs is to co-apply biosolids with biochar to soil, because biochar's chemical (e.g., aromaticity) and physical properties (e.g., surface area) give it a high affinity to bind many organic chemicals in the environment. We developed a soil-specific extraction method and utilized a luciferase-based bioassay (CALUX) to detect endocrine active chemicals in a biosolids-biochar co-amendment soil greenhouse study. Both biochar (walnut shell, 900 °C) and biosolids had positive impacts on carrot and lettuce biomass accumulation over our study period. However, the walnut shell biochar stimulated aryl hydrocarbon receptor activity, suggesting the presence of potential endocrine active chemicals in the biochar. Since the biochar rate tested (100 t ha<sup>-1</sup>) is above the average agronomic rate (10–20 t ha<sup>-1</sup>), endocrine effects would not be expected in most environmental applications. The effect of high temperature biochars on endocrine system pathways must be explored further, using both quantitative analytical tools to identify potential endocrine active chemicals and highly sensitive bioanalytical assays such as CALUX to measure the resulting biological activity of such compounds.

© 2017 Elsevier Ltd. All rights reserved.

\* Corresponding author.

E-mail address: [sjparikh@ucdavis.edu](mailto:sjparikh@ucdavis.edu) (S.J. Parikh).

## 1. Introduction

Biosolids are the repository for a substantial portion of nutrients consumed in the form of food. Returning biosolids to food production systems can help “close the loop” in nutrient cycling (DeLuca, 2009). A challenge is that biosolids contain not only plant nutrients, but also pharmaceuticals (Table A.1) (U.S. Environmental Protection Agency, 2009).

Many organic contaminants commonly found in biosolids, such as pharmaceuticals and personal care products, sorb strongly to organic matter (Aristilde and Sposito, 2010; Carmosini and Lee, 2009; Rogers, 1996) and clays (Carmosini and Lee, 2009; Carrasquillo et al., 2008) due in part to their low solubility and the interactive effects of their multiple functional groups (MacKay and Seremet, 2008). Many pharmaceuticals exhibit a high sorption affinity to biosolids (Wu et al., 2009). For example, the antibiotic ciprofloxacin is not significantly removed after methanogenic wastewater treatment due to its high sorption properties (Golet et al., 2003). The antibacterials triclosan (TCS) and triclocarban (TCC) have low water solubility, high octanol-water partitioning coefficients ( $K_{ow}$ ), and high organic carbon-water partitioning coefficients ( $K_{oc}$ ) (Table A.2), and thus have a moderate affinity for solids. These compounds accumulate in biosolids and are found in nearly all wastewater treatment plant (WWTP) effluents (U.S. Environmental Protection Agency, 2009). Additionally, biodegradation studies have demonstrated that some pharmaceuticals or their residues can persist in the soil environment for over six months after application (Boxall et al., 2006). Together, these properties suggest that pharmaceuticals applied to land in biosolids could accumulate and persist in the environment (Näslund et al., 2008).

Ciprofloxacin, TCS, and TCC are classified as “contaminants of emerging concern” by the US EPA, a designation for substances present at trace but potentially ecotoxicologically critical concentrations (Lozano et al., 2010). While many pharmaceutical residues in environmental samples are relatively low in concentration, especially compared to the prescribed human dosage, non-specific bioactivity of these drugs might reach environmentally-relevant levels, causing adverse effects on soil biota and biologically-driven ecosystem services (Dolliver et al., 2007; Jones-Lepp et al., 2010; Kumar et al., 2005; Lillenberg et al., 2010). It is of particular concern that several emerging contaminants, including TCS (Foran et al., 2000) and TCC (Chen et al., 2008) have capacity for endocrine-disrupting activity, or disruption of normal hormonal functions in biota. Since land application is a common means for disposal of biosolids produced at US wastewater treatment facilities (National Research Council, 2002), cost-effective technologies are needed to prevent associated pharmaceuticals from contaminating the environment or affecting human health.

Biochar, a carbonized byproduct of the pyrolysis of organic biomass, is an increasingly common soil amendment and may provide agronomic benefits (Laird et al., 2010; Lehmann et al., 2003; Van Zwieten et al., 2010). Using biochar as a co-amendment with biosolids could potentially reduce bioavailability and toxicity of endocrine-disrupting chemicals (EDCs) found in biosolids. Because of its high aromaticity and surface area, biochar can effectively sorb many organic chemicals introduced into the environment (Jung et al., 2013; Yao et al., 2012). This, in turn, could decrease bioavailability and limit bioaccumulation of EDCs in plants (Khan et al., 2013; Yu et al., 2009). While use of biochar has been investigated in treating heavy metals (Park et al., 2011) and pesticides (Yu et al., 2009), the effect of biochar co-amendment with biosolids on the bioavailability and resulting activity of EDCs is unknown. The physical-chemical properties of biochar (e.g., pH, surface area, elemental composition) vary depending on

parameters such as feedstock and pyrolysis temperature (Mukome et al., 2013b), and biochars produced via gasification have been shown to have high levels of EDCs such as polycyclic aromatic hydrocarbons (PAHs) (Cole et al., 2012; Hale et al., 2012).

Pharmaceutical residues in environmental samples are often analyzed using chemical methods such as liquid chromatography tandem mass spectrometry (LC/MS/MS), which can resolve trace environmental concentrations of specific compounds of known mass with high precision. However, such analyses measure a total concentration and not the concentration of the contaminant that will actually be biologically available.

An alternative approach is application of the Chemically-Activated Luciferase Expression (CALUX) bioassay (Denison et al., 2004) that can measure the combined ability of chemicals extracted from environmental matrices, including biosolids- and biochar-amended soil, to stimulate hormone receptor signaling pathways. These CALUX bioassays are recombinant cell lines that contain a stably transfected nuclear receptor responsive firefly luciferase reporter gene and they respond to activators/inhibitors of these receptors in a chemical-, time-, concentration- and receptor-specific manner with the induction/inhibition of luciferase activity and measurement of light output (Denison et al., 2004). CALUX bioassays based on estrogen receptor (ER), androgen receptor (AR), glucocorticoid receptor (GR), progesterone receptor (PR) and aryl hydrocarbon receptor (AhR) have been previously developed and used to determine the total biological activity (activation/inhibition) of a chemical(s) or extract on a given receptor pathway (Giudice and Young, 2011; Lorenzen et al., 2004; Topp et al., 2006; Zhang et al., 2009). Measurement of endocrine active chemicals in biosolids with CALUX bioassays have been used as a method both to characterize such activity (Lorenzen et al., 2004; Topp et al., 2006; Zhang et al., 2009) and for environmental mobilization studies (Giudice and Young, 2011). In addition to showing high sensitivity and specificity for known endocrine-disrupting hormone receptor agonists and antagonists (van der Linden et al., 2008), the CALUX bioassay can provide an estimate of the overall activity of mixtures of compounds such as may be found in biosolids or in soil after biosolids application.

In this study, we measured the individual and combined effects of biochar and biosolids co-amendments on endocrine active chemicals in soil from greenhouse-grown lettuce and carrots. We utilized CALUX bioassays to specifically target androgen, estrogen, aryl hydrocarbon, and glucocorticoid/progesterone receptors, and hypothesized that biochar preferentially sorbs organic pharmaceuticals decreasing their bioavailability in the soil and thus potential to produce endocrine disrupting effects. Therefore, we expected to see lower bioassay activity levels in the biosolids-biochar treatments with high biochar application rates.

## 2. Materials and methods

### 2.1. Soil, biosolids, and biochar

A greenhouse experiment was conducted using a loamy sand (Columbia series, Aquic Xerochlovent) collected from USDA-NRCS Lockeford Plant Materials Center (Lockeford, CA). Soil particle size distribution was determined using the hydrometer method (82.5% sand, 6.0% clay) and water-holding capacity was determined using a common field method whereby soils were saturated and allowed to drain until drainage stopped ( $0.52 \text{ g g}^{-1}$ ). Biosolids were provided by the Woodland Wastewater Treatment Plant (Woodland, CA), derived from an extended aeration activated sludge process. Macronutrient (N, P, K) data for soil and biosolids were determined by the combustion method for total N (AOAC Official Method 972.43), a KCl extraction using a flow injection analyzer for  $\text{NO}_3\text{-N}$

and  $\text{NH}_4\text{-N}$  (Table A.3) (Knepel, 2003), the Olsen-P method for bioavailable P (Prokopy, 1995), and ammonium acetate displacement for exchangeable K (Thomas, 1982). The application rate of biosolids used was  $63.1 \text{ g biosolids kg}^{-1}$  soil based on plant demand for nitrogen, calculated at a rate of  $200 \text{ kg N ha}^{-1}$  to 30 cm (Evanylo et al., 2006). Non biosolids-amended treatments received the equivalent N in mineral fertilizer. Based on the 16-16-16 fertilizer (Lilly Miller Brands, Walnut Creek, CA, USA) used,  $87.4 \text{ kg ha}^{-1}$  phosphorus,  $166 \text{ kg ha}^{-1}$  potassium and  $62.5 \text{ kg ha}^{-1}$  sulfur were also applied. The biochar was made from walnut shells (*Juglans californica*) using a Biomax 50 downdraft gasifier at Dixon Ridge Farms (Winters, CA, USA) (Mukome et al., 2013a), as part of a waste-to-energy process (Pujol Pereira et al., 2016). Detailed information on the physical and chemical characteristics of this biochar have been detailed elsewhere (Mukome et al., 2013b). The pyrolysis temperature was  $900^\circ\text{C}$ , resulting in an ash content of 46.6% and a specific surface area of  $227.1 \text{ m}^2 \text{ g}^{-1}$  and total N of 0.47% (Mukome et al., 2013b). The walnut shell biochar was chosen specifically because it has been shown to perform well compared to other chars as a sorbent for pharmaceutical (and other) contaminants (Bair et al., 2016).

## 2.2. Chemicals

Biosolids were spiked with ciprofloxacin, TCS, and TCC (Sigma-Aldrich, St. Louis, MO, USA), at concentrations slightly higher than the maximum concentrations found in municipal biosolids from throughout the US (Table A.1) (U.S. Environmental Protection Agency, 2009). The spiking levels were determined by a mass balance approach to estimate the amount of plant-available pharmaceuticals based on sorption coefficients of the soil, biosolids, and biochar. The pharmaceuticals were spiked in air-dried biosolids in 0.1% formic acid in methanol, at concentrations of  $100 \text{ mg kg}^{-1}$  ciprofloxacin,  $200 \text{ mg kg}^{-1}$  TCS, and  $500 \text{ mg kg}^{-1}$  TCC. The methanol was allowed to evaporate in a fume hood for 24 h before application of the biosolids to soil.

## 2.3. Experimental design and sample collection

A leaf crop (lettuce, *Lactuca sativa* L.) and root crop (carrot, *Daucus carota*) were planted in 4-L pots of 2 mm-sieved soil in a factorial design that combined three levels of biochar application (0, 10, and  $100 \text{ t ha}^{-1}$ ), and two levels of biosolids application (0 and  $20 \text{ t ha}^{-1}$ ). A high biochar application ( $100 \text{ t ha}^{-1}$ ) was chosen to assess if this rate resulted in more benefits (e.g., increased plant growth or decreased pharmaceutical bioavailability) than a more standard application rate ( $10 \text{ t ha}^{-1}$ ). A high biochar application rate ( $100 \text{ t ha}^{-1}$ ) has been shown to maximize crop productivity (Jeffery et al., 2011), and while this rate may not be economically feasible in broadcast application, targeted (banded) application may allow for high targeted rates of biochar inputs (Blackwell et al., 2010). Each biosolids-biochar treatment consisted of five replicates per plant type. Water-holding capacity was determined as detailed above for each biosolids-biochar treatment, and was maintained across treatments at 60–70% with DI water. Plants were grown in a temperature- and light-controlled greenhouse (Hoagland Hall, UC Davis, CA) from November 30, 2012, until harvest on January 30, 2013 (62 d). Upon harvest, soil samples were taken and deep-frozen ( $-80^\circ\text{C}$ ) until further analysis. Above- and below-ground plant biomass values were determined after freeze-drying plant samples for 48 h.

## 2.4. Sample preparation and CALUX analysis

To directly assess the bioavailability of these compounds from

soil, all samples were extracted with  $18.2 \text{ M}\Omega\text{-cm}$  water (Barnstead Nanopure) instead of organic solvents (Andersson et al., 2009; Hale et al., 2012; Kanematsu et al., 2009). All references to the water extraction are to  $18.2 \text{ M}\Omega\text{-cm}$  water. All extraction glassware was washed, rinsed in water, acid-washed in  $1 \text{ N H}_2\text{SO}_4$  and autoclaved for 30 min. Two grams of 2 mm-sieved air-dried soil in glass centrifuge tubes with Teflon-lined caps was extracted with 30 mL water in an ultrasonication bath for 30 min in the dark, followed by centrifugation at 3500 rpm for 20 min. The supernatant was filtered through Whatman GF/F glass fiber filters using Buchner funnels into glass culture tubes and the filtrate acidified to pH 2 with HCl. The sample was then concentrated using solid phase Oasis HLB 6-cc cartridges (Waters Corporation, Milford, MA) (Giudice and Young, 2011), and evaporated to dryness in glass vials at room temperature under a gentle stream of nitrogen. The dried residues were resuspended in  $50 \mu\text{L}$  of DMSO and analyzed using CALUX bioassays to determine the presence and relative concentration of activators of ER, AR, GR, PR and AhR signaling pathways (ER:  $17\beta$  Estradiol, AR: dihydroxytestosterone, GR: dexamethasone, PR: progesterone, and AhR: 2,3,7,8-tetrachlorodibenzo-p-dioxin (TCDD)). CALUX bioassays were carried out as previously described (Giudice and Young, 2011) and the results expressed relative to the activity obtained with a maximal inducing concentration of the positive control chemicals for each receptor-selective CALUX bioassay, normalized per gram dry soil to allow for cross-treatment comparison.

## 2.5. LC/MS/MS

HPLC-ESI/MS/MS was used to determine recovery efficiency of the extraction method (SI, Tables A.4–6).

## 2.6. Statistical analysis

All statistical analyses were performed using the R computing package (R Core Team, 2016) version 3.2.4. For analysis of the CALUX bioassay, data from the lettuce and carrot trials were pooled. Two-way analysis of variance (ANOVA) was performed for CALUX bioassay and plant biomass data with biosolids and biochar levels as factors. We found no significant effect of the biosolids  $\times$  biochar interaction and therefore proceeded with analysis with biosolids and biochar as main effects. Tukey's honest significant differences (HSD) were used as post hoc tests to assess treatment differences. Differences were determined to be statistically significant if  $p < 0.05$ . The relationship between CALUX bioassay activity levels and plant growth were assessed using regression analysis.


## 2.7. Data and analysis code availability

All data and analysis code will be available at <https://github.com/carolynanderson/calux>.

# 3. Results and discussion

## 3.1. CALUX bioassay detects endocrine active chemicals in biosolids-biochar co-amended soil

Soil samples from both the lettuce and carrot trials show low levels of AhR activity, in all combinations of biosolids-biochar treatments (Fig. 1, Table 1). The addition of biosolids significantly increased the AhR activity in soils compared to no biosolids treatments (mean values from 4.7% to 6.9%), as did the addition of biochar compared to no biochar treatments (mean values from 4.3% to 7.4%), but this was only significant at the highest biochar application rate ( $100 \text{ t ha}^{-1}$ ). These results suggest that biosolids and



**Fig. 1.** Aryl hydrocarbon receptor (AhR) activity in pooled soil extracts from biosolids-biochar treatments, as measured by the AhR-CALUX bioassay. AhR activity as expressed as relative light units normalized to the activity observed with a maximally inducing concentration of TCDD. Box plots show medians with the middle bars, surrounded on the top and bottom by the first and third quartiles, respectively. The highest and lowest data points within 1.5 times the interquartile range are shown by the whiskers; outliers are represented as points.

biochar contain AhR active chemicals; this is supported by previous studies reporting the presence of the AhR agonist TCS in biosolids in trace quantities (Ahn et al., 2008). Interestingly, extracts from soil samples revealed no activators of the ER, AR, GR or PR CALUX bioassays (data not shown); the presence of antagonists of these receptors was not determined.

More specifically, in non-biosolids amended treatments, AhR activity increased with increasing biochar application rate (Fig. 1), suggesting that this 900 °C walnut shell biochar itself may be a source for AhR active compounds. Based on this result, an extensive non-target analysis for biochar-associated compounds which may responsible for increased AhR activity is warranted. However, some of the compounds in biochar that could increase AhR activity include PAHs and dioxins (Quilliam et al., 2013; Stevens et al., 2009), both byproducts of biochar production, depending on the feedstock and production process and temperature (Hajaligol et al., 2001; Hale et al., 2012). Specifically, PAH levels exceeding environmental standards have been detected in biochars produced via gasification (i.e., in the presence of oxygen) (Cole et al., 2012; Hale

et al., 2012), the method used to form the walnut shell biochar used in this study. This is in contrast to biochars formed via pyrolysis (i.e., in oxygen-limited conditions), which show decreasing PAH concentrations with increasing pyrolysis temperatures (Hale et al., 2012; Keiluweit et al., 2012; Zielińska and Oleszczuk, 2016). It is important to note that while the highest biochar application rate tested in our study ( $100 \text{ t ha}^{-1}$ ) was over 10 times the most common agronomic rate used in land application, benefits (as measured by crop productivity) have been demonstrated with biochar applications at this rate (Glaser et al., 2002; Jeffery et al., 2011). However, high rates of biochar application have also been shown to cause a change in microbial community composition and a physiological response to stress in Gram-negative bacteria, in addition to a significant reduction in soil nitrate (Ippolito et al., 2014). Additionally, our use of an environmentally-relevant water-based extraction method most likely contributed to low concentrations of PAHs and dioxins extracted from these samples compared to that which could be obtained with a nonpolar solvent extraction method (e.g. methanol, toluene) (Andersson et al., 2009; Hale et al., 2012; Kanematsu et al., 2009), suggesting that endocrine active compounds may not be of significant concern under environmental conditions.

The relationship between individual biochar feedstock properties and the concentration and composition of PAHs and other AhR ligands is still poorly understood, as is the environmental fate of PAHs in a soil application of biochar (Bucheli et al., 2015; Dutta et al., 2016). Given that the AhR can bind a wide range of structurally diverse chemicals (DeGroot et al., 2011; Denison et al., 2011), biochar and biosolids could contain a variety of active chemicals, the identity of which remains to be determined. This warrants more research to assess AhR activity-inducing compounds in different biochars, and determine whether the presence of these compounds has negative implications for environmental or human health. The CALUX method is a novel approach that can directly and sensitively detect such activity, and can be used in tandem with traditional analytical approaches which by themselves may not be sufficient (Mayer et al., 2016).

### 3.2. Biochar and biosolids effects on lettuce growth

Only biochar, not biosolids, significantly affected lettuce shoot growth. The presence of biochar significantly increased shoot growth ( $10 \text{ t ha}^{-1}$  biochar =  $16.0 \pm 0.32 \text{ g}$ ;  $100 \text{ t ha}^{-1}$  biochar =  $16.07 \pm 0.48 \text{ g}$ ) compared to no biochar ( $15.22 \pm 0.69 \text{ g}$ ; Fig. 2a). In non-biosolids amended treatments, the presence of biochar significantly increased lettuce shoot growth ( $16.15 \pm 0.33 \text{ g}$  for  $10 \text{ t ha}^{-1}$  biochar,  $15.91 \pm 0.38 \text{ g}$  for  $100 \text{ t ha}^{-1}$  biochar) compared to the non-biochar treatment ( $15.09 \pm 0.34 \text{ g}$ ).


Both biochar and biosolids treatments had significant effects on lettuce root growth. Lettuce plants that received any level of biochar had significantly higher root biomass ( $10 \text{ t ha}^{-1}$  biochar =  $12.96 \pm 0.17 \text{ g}$ ;  $100 \text{ t ha}^{-1}$  biochar =  $13.12 \pm 0.25 \text{ g}$ ), than those without biochar ( $12.60 \pm 0.16 \text{ g}$ ) (Fig. 2b). Biochar has been shown to stimulate root growth by increasing water and nutrient availability, aeration, and soil pH (Lehmann et al., 2011) (although in the current study plants were watered regularly, and thus positive effects on water retention are probably negligible).

**Table 1**

Two-way ANOVA of AhR activity levels, with biosolids and biochar application rates as factors. Differences were determined to be statistically significant if  $p < 0.05$ .

	Degrees of freedom, d.f.	Sum of squares, SS	Mean squares, MS	F-statistic	p-value
Biosolids	1	75.31	75.31	6.13	0.0163
Biochar	2	123.18	61.59	5.02	0.0099
Residuals	56	687.68	12.28		





**Fig. 2.** Growth differences in lettuce and carrot shoots (a) and roots (b) in biosolids and biochar treatments. Box plots show medians with the middle bars, surrounded on the top and bottom by the first and third quartiles, respectively. The highest and lowest data points within 1.5 times the interquartile range are shown by the whiskers; outliers are represented as points.

Biochar also appears to mitigate potentially negative effects of biosolids on lettuce root growth, but only at the highest biochar application rate in this study (Fig. 2). In treatments without biosolids, lettuce root biomass significantly increased with the addition of biochar ( $10 \text{ t ha}^{-1}$  biochar =  $13.21 \pm 0.11$ ;  $100 \text{ t ha}^{-1}$  biochar =  $13.24 \pm 0.22$ ) compared to no biochar ( $12.63 \pm 0.23 \text{ g}$ ). In the biosolids treatments, biochar significantly increased lettuce root biomass only in the  $100 \text{ t ha}^{-1}$  biochar treatment ( $13.00 \pm 0.23 \text{ g}$ ) compared to the no biochar treatment ( $12.57 \pm 0.07 \text{ g}$ ). Further, in the absence of biochar there were no significant differences in lettuce root growth in biosolids-amended and unamended treatments. It is possible that compounds in biosolids (e.g., salts, excess nutrients) might negatively affect lettuce root growth (Andrés et al., 2011), and these results suggest that a high biochar application rates can overcome this. Similar greenhouse studies show that biochar feedstock type is an important determinant in the plant response to biochar. Different biochar feedstocks have been shown to increase corn, grass, and barley growth (Griffin et al., 2017; Jones et al., 2012; Prendergast-Miller et al., 2014; Rajkovich et al., 2012), although contrary to the results of the current study, this effect can be negated at high biochar application rates (e.g.,  $91 \text{ t ha}^{-1}$ ) (Rajkovich et al., 2012). Further, the effect of biochar can differ in greenhouse vs. field studies (Jones et al., 2012; Rajkovich et al., 2012).

Together, these root and shoot growth data suggest that the walnut shell biochar has a greater impact on lettuce growth than do biosolids, at least at the given application rates. While biochar on its own has been shown to increase plant growth (Khan et al., 2013;

Lehmann et al., 2003), this positive effect could be negated by toxicity from high levels of volatile organic matter in some biochars (Deenik et al., 2010; Lehmann et al., 2011). Furthermore, while biosolids have been shown to increase crop yield (Binder et al., 2002; Christie et al., 2001; Singh and Agrawal, 2008), potential differences in lettuce biomass in the current study between biosolids and non-biosolids treatments might have been obscured by additions of N fertilizer to non-biosolids pots to remove differences in fertility among the treatments.


### 3.3. Biochar and biosolids effects on carrot growth

In the carrot treatments, there were no significant differences between biosolids and biochar treatments in root or shoot growth (Fig. 2). In contrast to what was observed in lettuce, the addition of biosolids, regardless of biochar concentration, significantly increased carrot shoot mass. This pattern of increased plant growth with biosolids was different from that found in the lettuce plants, suggesting that there might be different growth responses between lettuce and carrot plants in the presence of compounds found in biosolids. Differences in contaminant uptake in leaf and root crops have been demonstrated and attributed to both physiological and chemical differences in the plants. For example, the mechanisms of uptake of veterinary medicines by lettuce and carrot plants have been shown to follow Gaussian and non-Gaussian relationships, respectively (Boxall et al., 2006). Differences in uptake have also been ascribed to the chemistry of the contaminant and its matrix; for example, low uptake of TCS in radish root has been attributed to low ionization of TCS in the study soil (Carter et al., 2014). Such plant growth differences warrant further research, but this is outside the scope of the current study. It may be that any negative effects of pharmaceuticals in biosolids on carrot root and shoot growth are mitigated by the high carbon and nutrient contents of biosolids. Such trends are seen in crop yields (Singh and Agrawal, 2008) and in microbially-driven soil nutrient cycling processes, which either increase or remain unaltered in the presence of biosolids (Park et al., 2013). It is also possible that pharmaceuticals or other contaminants in biosolids are sorbed strongly to the organic matter matrix of the biosolid itself, or biochar if present, limiting plant uptake (Yao et al., 2012), although the sorption dynamics depend on the individual contaminant and biosolids or biochar matrix. Further, as previously mentioned, the uptake dynamics between lettuce versus carrot plants depend on physiological differences as well as chemical differences in the compound, which may explain why these plants responded differently to the treatments in regards to growth.

### 3.4. Environmental implications

While we hypothesized that the addition of biochar would decrease endocrine activity associated with biosolids treatments, high biochar application rates ( $100 \text{ t ha}^{-1}$ ) actually increased activity levels, albeit by a small but significant amount. This suggested that this particular biochar may be a source of AhR active compounds, which may reflect the presence of PAHs or numerous other classes of AhR agonists (DeGroot et al., 2011; Denison et al., 2011) in the biochar-amended soil. Together, these results indicate that while biochar (walnut shell,  $900^\circ\text{C}$ ) can positively affect plant growth, it might also contribute chemicals into the soil that potentially could have endocrine-disrupting activity. The positive CALUX bioassay results presented here suggest the need for non-target analysis of the biochar to confirm and quantify which chemicals are inducing this response.

The biochar rate that induced this effect ( $100 \text{ t ha}^{-1}$ ) was above the average agronomic rate ( $10\text{--}20 \text{ t ha}^{-1}$ ), so while a small



**Fig. 3.** Regressions of plant biomass versus soil AhR activity for (a) shoots and (b) roots. P-values from linear regressions are reported in each panel.

increase in AhR activity was observed with the 10 t ha<sup>-1</sup> sample, it is unlikely that biochar at normal environmental application levels would be of toxicological concern. Further, the AhR activity in the soil does not appear to be coupled with effects on plant growth (Fig. 3,  $R^2 < 0.10$  and  $p > 0.10$  for all plant/tissue combinations), suggesting that low AhR activity associated with biochar does not affect crop yield.

The PAHs potentially present in biochars may be taken up by crops, resulting in a potential source of exposure to humans (Zohair et al., 2006). Therefore, to ensure agronomic safety of biochar application, the presence and activity of endocrine-active compounds in biochar should be explored further, using both quantitative analytical tools to measure PAHs, dioxins, and other endocrine-active compounds. Highly sensitive bioanalytical assays such as CALUX can help to assess the biological activity of such compounds. While the biochar used in this study (walnut shell, 900 °C) is an isolated example and these results do not apply to all chars, the method developed in this study provides a tool for rapid screening of complex environmental matrices including other biochars, and the results suggest a need to perform non-target analysis on this char. CALUX has considerable potential as a low-cost, sensitive tool to screen biological and ecological effects of complex mixtures of compounds in environmental samples.

## Acknowledgments

This research was supported by the UC Davis Superfund Trainee Program (funded through the National Institute of Environmental Health Sciences of the National Institutes of Health under Award Number P42ES004699) and the Soils and Biogeochemistry Graduate Group. We thank the Lockeford Plant Materials Center and the Woodland Wastewater Treatment Plant for supplying the soil and biosolids used in this study. S.J. Parikh acknowledges support from the United States Department of Agriculture (USDA) National Institute of Food and Agriculture (NIFA) for Hatch and multisite regional project (W-2082) funding. This work is/was supported by the USDA National Institute of Food and Agriculture, Hatch Project

CA-2122-H. The content is solely the responsibility of the authors and does not necessarily represent the official views of the National Institutes of Health, USDA, or other funding sources.

## Appendix A. Supplementary data

Supplementary data related to this article can be found at <http://dx.doi.org/10.1016/j.chemosphere.2017.04.035>.

## References

- Ahn, K.C., Zhao, B., Chen, J., Cherednichenko, G., Sanmarti, E., Denison, M.S., Lasley, B., Pessah, I.N., Kultz, D., Chang, D.P.Y., Gee, S.J., Hammock, B.D., 2008. In vitro biologic activities of the antimicrobials triclocarban, its analogs, and triclosan in bioassay screens: receptor-based bioassay screens. *Environ. Health Perspect.* 116, 1203–1210. <http://dx.doi.org/10.1289/ehp.11200>.
- Andersson, E., Rotander, A., von Kronhelm, T., Berggren, A., Ivarsson, P., Hollert, H., Engwall, M., 2009. AhR agonist and genotoxicant bioavailability in a PAH-contaminated soil undergoing biological treatment. *Environ. Sci. Pollut. Res.* 16, 521–530. <http://dx.doi.org/10.1007/s11356-009-0121-9>.
- Andrés, P., Mateos, E., Tarrasón, D., Cabrera, C., Figuerola, B., 2011. Effects of digested, composted, and thermally dried sewage sludge on soil microbiota and meso-fauna. *Appl. Soil Ecol.* 48, 236–242. <http://dx.doi.org/10.1016/j.apsoil.2011.03.001>.
- Aristilde, L., Sposito, G., 2010. Binding of ciprofloxacin by humic substances: a molecular dynamics study. *Environ. Toxicol. Chem.* 29, 90–98. <http://dx.doi.org/10.1002/etc.19>.
- Bair, D.A., Mukome, F.N.D., Popova, I.E., Ogunyoku, T.A., Jefferson, A., Wang, D., Hafner, S.C., Young, T.M., Parikh, S.J., 2016. Sorption of pharmaceuticals, heavy metals, and herbicides to biochar in the presence of biosolids. *J. Environ. Qual.* 45, 1998–2006. <http://dx.doi.org/10.2134/jeq2016.03.0106>.
- Binder, D.L., Dobermann, A., Sander, D.H., Cassman, K.G., 2002. Biosolids as nitrogen source for irrigated maize and rainfed sorghum. *Soil Sci. Soc. Am. J.* 66, 531–543.
- Blackwell, P., Krull, E., Butler, G., Herbert, A., Solaiman, Z., 2010. Effect of banded biochar on dryland wheat production and fertiliser use in south-western Australia: an agronomic and economic perspective. *Aust. J. Soil Res.* 48, 531. <http://dx.doi.org/10.1071/SR10014>.
- Boxall, A.B.A., Johnson, P., Smith, E.J., Sinclair, C.J., Stutt, E., Levy, L.S., 2006. Uptake of veterinary medicines from soils into plants. *J. Agric. Food Chem.* 54, 2288–2297. <http://dx.doi.org/10.1021/jf053041t>.
- Bucheli, T., Hilber, I., Schmidt, H., 2015. Polycyclic aromatic hydrocarbons and polychlorinated aromatic compounds in biochar. In: Lehmann, J., Joseph, S. (Eds.), *Biochar for Environmental Management: Science, Technology and Implementation*. Earthscan, London, UK, pp. 595–624.
- Carmosini, N., Lee, L.S., 2009. Ciprofloxacin sorption by dissolved organic carbon from reference and bio-waste materials. *Chemosphere* 77, 813–820. <http://dx.doi.org/10.1016/j.chemosphere.2009.08.003>.
- Carrasquillo, A.J., Bruland, G.L., MacKay, A.A., Vasudevan, D., 2008. Sorption of ciprofloxacin and oxytetracycline zwitterions to soils and soil minerals: influence of compound structure. *Environ. Sci. Technol.* 42, 7634–7642.
- Carter, L.J., Harris, E., Williams, M., Ryan, J.J., Kookana, R.S., Boxall, A.B.A., 2014. Fate and uptake of pharmaceuticals in soil–plant systems. *J. Agric. Food Chem.* 62, 816–825. <http://dx.doi.org/10.1021/jf404282y>.
- Chen, J., Ahn, K.C., Gee, N.A., Ahmed, M.I., Duleba, A.J., Zhao, L., Gee, S.J., Hammock, B.D., Lasley, B.L., 2008. Triclocarban enhances testosterone action: a new type of endocrine disruptor? *Endocrinology* 149, 1173–1179. <http://dx.doi.org/10.1210/en.2007-1057>.
- Christie, P., Easson, D.L., Picton, J.R., Love, S.C., 2001. Agronomic value of alkaline-stabilized sewage biosolids for spring barley. *Agron. J.* 93, 144–151.
- Cole, D.P., Smith, E.A., Lee, Y.J., 2012. High-resolution mass spectrometric characterization of molecules on biochar from pyrolysis and gasification of switchgrass. *Energy Fuels* 26, 3803–3809. <http://dx.doi.org/10.1021/ef300356u>.
- Deenik, J.L., McClellan, T., Uehara, G., Antal, M.J., Campbell, S., 2010. Charcoal volatile matter content influences plant growth and soil nitrogen transformations. *Soil Sci. Soc. Am. J.* 74, 1259. <http://dx.doi.org/10.2136/sssaj2009.0115>.
- DeGroot, D., He, G., Fracalvieri, D., Bonati, L., Pandini, A., Denison, M.S., 2011. AhR ligands: promiscuity in binding and diversity in response. In: *The Ah Receptor in Biology and Toxicology*. John Wiley and Sons, Hoboken, NJ, pp. 63–79.
- DeLuca, T.H., 2009. Nutrient imbalances: follow the waste. *Science* 326, 665.
- Denison, M.S., Soshilov, A.A., He, G., DeGroot, D.E., Zhao, B., 2011. Exactly the same but different: promiscuity and diversity in the molecular mechanisms of action of the aryl hydrocarbon (dioxin) receptor. *Toxicol. Sci.* 124, 1–22. <http://dx.doi.org/10.1093/toxsci/kfr218>.
- Denison, M.S., Zhao, B., Baston, D.S., Clark, G.C., Murata, H., Han, D., 2004. Recombinant cell bioassay systems for the detection and relative quantitation of halogenated dioxins and related chemicals. *Talanta* 63, 1123–1133. <http://dx.doi.org/10.1016/j.talanta.2004.05.032>.
- Dolliver, H., Kumar, K., Gupta, S., 2007. Sulfamethazine uptake by plants from manure-amended soil. *J. Environ. Qual.* 36, 1224. <http://dx.doi.org/10.2134/jeq2006.0266>.

- Dutta, T., Kwon, E., Bhattacharya, S.S., Jeon, B.H., Deep, A., Minori, U., Kim, K.-H., 2016. Polycyclic aromatic hydrocarbons (PAHs) and volatile organic compounds (VOCs) in biochar and biochar-amended soil: a review. *GCB Bioenergy*. <http://dx.doi.org/10.1111/gcbb.12363> n/a–n/a.
- Evanylo, Greg K., Abaye, A.O., Basden, T.J., Beegle, D., Binford, G.D., Daniels, W.L., Duiker, S.W., Evanylo, Gregory K., Haering, K.C., Hansen, D.J., Mullins, G., Taylor, R.W., 2006. The Mid-Atlantic Nutrient Management Handbook.
- Foran, C., Bennett, E., Benson, W., 2000. Developmental evaluation of a potential non-steroidal estrogen: triclosan. *Mar. Environ. Res.* 50, 153–156. [http://dx.doi.org/10.1016/S0141-1136\(00\)00080-5](http://dx.doi.org/10.1016/S0141-1136(00)00080-5).
- Giudice, B.D., Young, T.M., 2011. Mobilization of endocrine-disrupting chemicals and estrogenic activity in simulated rainfall runoff from land-applied biosolids. *Environ. Toxicol. Chem.* 30, 2220–2228. <http://dx.doi.org/10.1002/etc.631>.
- Glaser, B., Lehmann, J., Zech, W., 2002. Ameliorating physical and chemical properties of highly weathered soils in the tropics with charcoal - a review. *Biol. Fertil. Soils* 35, 219–230. <http://dx.doi.org/10.1007/s00374-002-0466-4>.
- Golet, E.M., Xifra, I., Siegrist, H., Alder, A.C., Giger, W., 2003. Environmental exposure assessment of fluoroquinolone antibacterial agents from sewage to soil. *Environ. Sci. Technol.* 37, 3243–3249. <http://dx.doi.org/10.1021/es0264448>.
- Griffin, D.E., Wang, D., Parikh, S.J., Scow, K.M., 2017. Short-lived effects of walnut shell biochar on soils and crop yields in a long-term field experiment. *Agric. Ecosyst. Environ.* 236, 21–29.
- Hajjalilol, M., Waymack, B., Kellogg, D., 2001. Low temperature formation of aromatic hydrocarbon from pyrolysis of cellulosic materials. *Fuel* 80, 1799–1807. [http://dx.doi.org/10.1016/S0016-2361\(01\)00063-1](http://dx.doi.org/10.1016/S0016-2361(01)00063-1).
- Hale, S.E., Lehmann, J., Rutherford, D., Zimmerman, A.R., Bachmann, R.T., Shitumbanuma, V., O'Toole, A., Sundqvist, K.L., Arp, H.P.H., Cornelissen, G., 2012. Quantifying the total and bioavailable polycyclic aromatic hydrocarbons and dioxins in biochars. *Environ. Sci. Technol.* 46, 2830–2838. <http://dx.doi.org/10.1021/es203984k>.
- Ippolito, J.A., Stromberger, M.E., Lentz, R.D., Dungan, R.S., 2014. Hardwood biochar influences calcareous soil physicochemical and microbiological status. *J. Environ. Qual.* 43, 681. <http://dx.doi.org/10.2134/jeq2013.08.0324>.
- Jeffery, S., Verheijen, F.G.A., van der Velde, M., Bastos, A.C., 2011. A quantitative review of the effects of biochar application to soils on crop productivity using meta-analysis. *Agric. Ecosyst. Environ.* 144, 175–187. <http://dx.doi.org/10.1016/j.agee.2011.08.015>.
- Jones, D.L., Rousk, J., Edwards-Jones, G., DeLuca, T.H., Murphy, D.V., 2012. Biochar-mediated changes in soil quality and plant growth in a three year field trial. *Soil Biol. Biochem.* 45, 113–124. <http://dx.doi.org/10.1016/j.soilbio.2011.10.012>.
- Jones-Lepp, T.L., Sanchez, C.A., Moy, T., Kazemi, R., 2010. Method development and application to determine potential plant uptake of antibiotics and other drugs in irrigated crop production systems. *J. Agric. Food Chem.* 58, 11568–11573. <http://dx.doi.org/10.1021/jf1028152>.
- Jung, C., Park, J., Lim, K.H., Park, S., Heo, J., Her, N., Oh, J., Yun, S., Yoon, Y., 2013. Adsorption of selected endocrine disrupting compounds and pharmaceuticals on activated biochars. *J. Hazard. Mater.* 263, 702–710. <http://dx.doi.org/10.1016/j.jhazmat.2013.10.033>.
- Kanematsu, M., Hayashi, A., Denison, M.S., Young, T.M., 2009. Characterization and potential environmental risks of leachate from shredded rubber mulches. *Chemosphere* 76, 952–958. <http://dx.doi.org/10.1016/j.chemosphere.2009.04.026>.
- Keilueit, M., Kleber, M., Sparrow, M.A., Simoneit, B.R.T., Prah, F.G., 2012. Solvent-extractable polycyclic aromatic hydrocarbons in biochar: influence of pyrolysis temperature and feedstock. *Environ. Sci. Technol.* 46, 9333–9341. <http://dx.doi.org/10.1021/es302125k>.
- Khan, S., Wang, N., Reid, B.J., Freddo, A., Cai, C., 2013. Reduced bioaccumulation of PAHs by *Lactuca sativa* L. grown in contaminated soil amended with sewage sludge and sewage sludge derived biochar. *Environ. Pollut.* 175, 64–68. <http://dx.doi.org/10.1016/j.envpol.2012.12.014>.
- Knepel, K., 2003. Determination of Nitrate in 2M KCl Soil Extracts by Flow Injection Analysis (No. QuikChem Method 12-107-04-1-B). Lachat Instruments, Loveland, CO.
- Kumar, K., Gupta, S.C., Baidoo, S.K., Chander, Y., Rosen, C.J., 2005. Antibiotic uptake by plants from soil fertilized with animal manure. *J. Environ. Qual.* 34, 2082. <http://dx.doi.org/10.2134/jeq2005.0026>.
- Laird, D.A., Fleming, P., Davis, D.D., Horton, R., Wang, B., Karlen, D.L., 2010. Impact of biochar amendments on the quality of a typical Midwestern agricultural soil. *Geoderma* 158, 443–449. <http://dx.doi.org/10.1016/j.geoderma.2010.05.013>.
- Lehmann, J., Kern, D., German, L., Mccann, J., Martins, G.C., Moreira, A., 2003. Soil fertility and production potential. In: *Amazonian Dark Earths*. Springer, pp. 105–124.
- Lehmann, J., Rillig, M.C., Thies, J., Masiello, C.A., Hockaday, W.C., Crowley, D., 2011. Biochar effects on soil biota – a review. *Soil Biol. Biochem.* 43, 1812–1836. <http://dx.doi.org/10.1016/j.soilbio.2011.04.022>.
- Lillenberg, M., Litvin, S.V., Nei, L., Roasto, M., Sepp, K., et al., 2010. Enrofloxacin and ciprofloxacin uptake by plants from soil. *Agron. Res.* 8, 807–814.
- Lorenzen, A., Hendel, J.G., Conn, K.L., Bittman, S., Kwabiah, A.B., Lazarovitz, G., Masse, D., McAllister, T.A., Topp, E., 2004. Survey of hormone activities in municipal biosolids and animal manures. *Environ. Toxicol.* 19, 216–225. <http://dx.doi.org/10.1002/tox.20014>.
- Lozano, N., Rice, C.P., Ramirez, M., Torrents, A., 2010. Fate of triclosan in agricultural soils after biosolid applications. *Chemosphere* 78, 760–766. <http://dx.doi.org/10.1016/j.chemosphere.2009.10.043>.
- MacKay, A.A., Seremet, D.E., 2008. Probe compounds to quantify cation exchange and complexation interactions of ciprofloxacin with soils. *Environ. Sci. Technol.* 42, 8270–8276. <http://dx.doi.org/10.1021/es800984x>.
- Mayer, P., Hilber, I., Gouliarmou, V., Hale, S.E., Cornelissen, G., Bucheli, T.D., 2016. How to determine the environmental exposure of PAHs originating from biochar. *Environ. Sci. Technol.* 50, 1941–1948. <http://dx.doi.org/10.1021/acs.est.5b05603>.
- Mukome, F.N.D., Six, J., Parikh, S.J., 2013a. The effects of walnut shell and wood feedstock biochar amendments on greenhouse gas emissions from a fertile soil. *Geoderma* 200–201, 90–98. <http://dx.doi.org/10.1016/j.geoderma.2013.02.004>.
- Mukome, F.N.D., Zhang, X., Silva, L.C.R., Six, J., Parikh, S.J., 2013b. Use of chemical and physical characteristics to investigate trends in biochar feedstocks. *J. Agric. Food Chem.* 61, 2196–2204. <http://dx.doi.org/10.1021/jf3049142>.
- Näslund, J., Hedman, J.E., Agestrand, C., 2008. Effects of the antibiotic ciprofloxacin on the bacterial community structure and degradation of pyrene in marine sediment. *Aquat. Toxicol.* 90, 223–227. <http://dx.doi.org/10.1016/j.aquatox.2008.09.002>.
- National Research Council, 2002. *Biosolids Applied to Land: Advancing Standards and Practices*. The National Academies Press, Washington, DC.
- Park, I., Zhang, N., Ogonyoku, T.A., Young, T.M., Scow, K.M., 2013. Effects of Triclosan and biosolids on microbial community composition in an agricultural soil. *Water Environ. Res. Res. Publ. Water Environ. Fed.* 85, 2237.
- Park, J.H., Choppala, G.K., Bolan, N.S., Chung, J.W., Chuasavathi, T., 2011. Biochar reduces the bioavailability and phytotoxicity of heavy metals. *Plant Soil* 348, 439–451. <http://dx.doi.org/10.1007/s11104-011-0948-y>.
- Prendergast-Miller, M.T., Duvall, M., Sohi, S.P., 2014. Biochar-root interactions are mediated by biochar nutrient content and impacts on soil nutrient availability: biochar-root interactions. *Eur. J. Soil Sci.* 65, 173–185. <http://dx.doi.org/10.1111/ejss.12079>.
- Prokopy, W., 1995. *Phosphorus in 0.5 M Sodium Bicarbonate Soil Extracts (No. QuikChem Method 12-115-01-1-B)*. Lachat Instruments, Milwaukee, WI.
- Pujol Pereira, E.I., Suddick, E.C., Six, J., 2016. Carbon abatement and emissions associated with the gasification of walnut shells for bioenergy and biochar production. *PLoS One* 11, e0150837. <http://dx.doi.org/10.1371/journal.pone.0150837>.
- Quilliam, R.S., Rangelcroft, S., Emmett, B.A., Deluca, T.H., Jones, D.L., 2013. Is biochar a source or sink for polycyclic aromatic hydrocarbon (PAH) compounds in agricultural soils? *GCB Bioenergy* 5, 96–103. <http://dx.doi.org/10.1111/gcbb.12007>.
- R Core Team, 2016. *R: a Language and Environment for Statistical Computing*. R Foundation for Statistical Computing, Vienna, Austria.
- Rajkovich, S., Enders, A., Hanley, K., Hyland, C., Zimmerman, A.R., Lehmann, J., 2012. Corn growth and nitrogen nutrition after additions of biochars with varying properties to a temperate soil. *Biol. Fertil. Soils* 48, 271–284. <http://dx.doi.org/10.1007/s00374-011-0624-7>.
- Rogers, H., 1996. Sources, behaviour and fate of organic contaminants during sewage treatment and in sewage sludge. *Sci. Total Environ.* 185, 3–26. [http://dx.doi.org/10.1016/0048-9697\(96\)05039-5](http://dx.doi.org/10.1016/0048-9697(96)05039-5).
- Singh, R.P., Agrawal, M., 2008. Potential benefits and risks of land application of sewage sludge. *Waste Manag.* 28, 347–358. <http://dx.doi.org/10.1016/j.wasman.2006.12.010>.
- Stevens, E.A., Mezrich, J.D., Bradfield, C.A., 2009. The aryl hydrocarbon receptor: a perspective on potential roles in the immune system. *Immunology* 127, 299–311. <http://dx.doi.org/10.1111/j.1365-2567.2009.03054.x>.
- Thomas, G.W., 1982. Exchangeable cations. In: Page, A.L. (Ed.), *Methods of Soil Analysis. Part 2. Chemical and Microbiological Properties*. American Society of Agronomy, Inc., Madison, WI, pp. 159–165.
- Topp, E., Burnison, K., Servos, M., Lorenzen, A., 2006. Persistence of endocrine-disrupting chemicals in agricultural soils. *J. Environ. Eng. Sci.* 5, 211–219. <http://dx.doi.org/10.1139/s05-016>.
- U.S. Environmental Protection Agency, 2009. *Targeted National Sewage Sludge Survey (No. EPA-822-R-08-018)* (Washington, DC).
- van der Linden, S.C., Heringa, M.B., Man, H.-Y., Sonneveld, E., Puijker, L.M., Brouwer, A., van der Burg, B., 2008. Detection of multiple hormonal activities in wastewater effluents and surface water, using a panel of steroid receptor CALUX bioassays. *Environ. Sci. Technol.* 42, 5814–5820. <http://dx.doi.org/10.1021/es702897y>.
- Van Zwieten, L., Kimber, S., Morris, S., Chan, K.Y., Downie, A., Rust, J., Joseph, S., Cowie, A., 2010. Effects of biochar from slow pyrolysis of papermill waste on agronomic performance and soil fertility. *Plant Soil* 327, 235–246. <http://dx.doi.org/10.1007/s11104-009-0050-x>.
- Wu, C., Sponberg, A.L., Witter, J.D., 2009. Sorption and biodegradation of selected antibiotics in biosolids. *J. Environ. Sci. Health Part A* 44, 454–461. <http://dx.doi.org/10.1080/10934520902719779>.
- Yao, Y., Gao, B., Chen, H., Jiang, L., Inyang, M., Zimmerman, A.R., Cao, X., Yang, L., Xue, Y., Li, H., 2012. Adsorption of sulfamethoxazole on biochar and its impact on reclaimed water irrigation. *J. Hazard. Mater.* 209–210, 408–413. <http://dx.doi.org/10.1016/j.jhazmat.2012.01.046>.
- Yu, X.-Y., Ying, G.-G., Kookana, R.S., 2009. Reduced plant uptake of pesticides with biochar additions to soil. *Chemosphere* 76, 665–671. <http://dx.doi.org/10.1016/j.chemosphere.2009.04.001>.
- Zhang, T., Yu, G., Wang, B., Fiedler, H., Huang, J., Deng, S., 2009. Bioanalytical characterization of dioxin-like activity in sewage sludge from Beijing, China.

- Chemosphere 75, 649–653. <http://dx.doi.org/10.1016/j.chemosphere.2008.12.064>.
- Zielińska, A., Oleszczuk, P., 2016. Effect of pyrolysis temperatures on freely dissolved polycyclic aromatic hydrocarbon (PAH) concentrations in sewage sludge-derived biochars. Chemosphere 153, 68–74. <http://dx.doi.org/10.1016/j.chemosphere.2016.02.118>.
- Zohair, A., Salim, A.-B., Soyibo, A.A., Beck, A.J., 2006. Residues of polycyclic aromatic hydrocarbons (PAHs), polychlorinated biphenyls (PCBs) and organochlorine pesticides in organically-farmed vegetables. Chemosphere 63, 541–553. <http://dx.doi.org/10.1016/j.chemosphere.2005.09.012>.